

Adding Explicit Solvent Molecules to Continuum Solvent Calculations for the Calculation of Aqueous Acid Dissociation Constants

Casey P. Kelly, Christopher J. Cramer, and Donald G. Truhlar**

Department of Chemistry and Supercomputing Institute, 207 Pleasant St. SE, University of Minnesota,
Minneapolis, MN 55455-0431

*cramer@chem.umn.edu, truhlar@chem.umn.edu

TITLE RUNNING HEAD: Calculation of Acid Dissociation Constants

Abstract: Aqueous acid dissociation free energies for a diverse set of 57 monoprotic acids have been calculated using a combination of experimental and calculated gas and liquid-phase free energies. For ionic species, aqueous solvation free energies were calculated using the recently developed SM6 continuum solvation model (Kelly, C. P.; Cramer, C. J.; Truhlar, D. G. *J. Chem. Theory Comput.* **2005**, *1*, 1133). This model combines a dielectric continuum with atomic surface tensions to account for bulk solvent effects. For some of the acids studied, a combined approach that involves attaching a single explicit water molecule to the conjugate base (anion), and then surrounding the resulting anion-water cluster by a dielectric continuum, significantly improves the agreement between the calculated pK_a value and experiment. This suggests that for some anions, particularly those concentrating charge on a single exposed heteroatom, augmenting implicit solvent calculations with a single explicit water molecule is required, and adequate, to account for strong short-range hydrogen bonding interactions between the anion and the solvent. Using the polyprotic acid H_2CO_3 , we also demonstrate the effect of adding several explicit waters by calculating the pK_a of bicarbonate (HCO_3^-) using as the conjugate base carbonate (CO_3^{2-}) bound by up to three explicit water molecules.

1. Introduction

Since Bjerrum's and Hammett's early work on substituent effects and solvent effects on acidity,¹ there has been continuing and substantial interest in predicting pK_a constants. Many of the methods that have been developed take advantage of empirical relationships between various molecular properties (either experimental or calculated) and pK_a , although recent work has focused on making pK_a predictions by directly calculating the free energy change associated with the acid dissociation process.²⁻⁴¹ This growing interest in direct pK_a calculation has occurred because empirical correlations between pK_a and various molecular properties have often been restricted to one or a few classes of compounds, such as substituted benzoic acids. Furthermore, in cases where correlations have been developed on the basis of larger and more diverse sets of compounds, it remains unclear whether these correlations can be readily applied to molecules outside the data sets within which they were originally developed, for example, their applicability to transition states, weak acids, and other compounds for which limited experimental pK_a data exist is open to question.

The most challenging aspect of directly calculating pK_a values is accurately accounting for the effect of solvent in a physical way. Solvation modeling is particularly challenging for ionic species due to strong electrostatic effects arising from unbalanced localized charges. For neutral species, implicit solvent models^{42,43} have been successful in predicting various liquid-phase properties. In these types of models, the electrostatic response of a bulk solvent is usually approximated as a homogeneous, dielectric continuum, making them an attractive alternative to computationally more demanding explicit solvent simulations. Explicit simulations require special treatment of finite-size effects⁴⁴ whereas implicit solvent models directly incorporate the bulk continuum limit. Implicit solvation models, though, require a decision as to which subsystem is to be treated as an explicit solute, with the remainder implicit. In a purely electrostatic implicit model, the implicit subsystem must be well-described by the bulk dielectric constant, but it is well-known that such a bulk model breaks down in the first solvation shell. Including implicit first-solvation-shell effects that depend on the atomic number⁴⁵ and geometry⁴⁶ of specific local solvent-exposed regions of the solute allows one to include these specific

solvation effects. Nevertheless questions remain about the minimum size of the system that must be treated explicitly to calculate quantitatively accurate free energies of solvation and pK_a constants. For example, for reactions involving RO^- , should one consider RO^- , $RO^-(H_2O)$, or $RO^-(H_2O)_n$ with $n > 1$ as the solute?

We recently developed an implicit solvation model for aqueous solvent, called Solvation Model 6 (SM6).⁴⁷ This model uses a dielectric continuum to treat electrostatic effects combined with atomic surface tensions to account for first-shell solvation effects, and it has been shown to give aqueous solvation free energies accurate to within ~ 0.5 kcal/mol for typical neutral solutes. During the development of this model, we found that some solutes containing concentrated regions of charge density (the majority of these being anions) required a parametrization significantly different from that of most other solutes in our training set. By replacing these solutes by their analogous monohydrated clusters (i.e., using a larger explicit region) we were able to obtain a single set of parameters that simultaneously fit (with reasonable accuracy) all of the neutral and ionic data in our training set. We concluded from this that strong local solute-solvent interactions play an important role in the aqueous solvation of some solutes, and in these cases, an implicit solvent model based on the minimal definition of the solute is not able to account for these interactions. Implicit solvation models that include one or more solvent molecules in the explicit subsystem have been called a combined dielectric-continuum model⁴⁸⁻⁵⁰ or a cluster-continuum model.^{51,52} We will also call it a combined implicit-explicit model.

Such a combined implicit-explicit solvent approach has seen recent use in direct pK_a calculations (some earlier applications focused on calculating solvatochromic shifts^{53,54}). Pliego and Riveros⁵² calculated pK_a values for 17 species using an implicit-explicit solvent approach for the charged species (for the neutral species, these workers did not include explicit water molecules) and were able to greatly improve the accuracy of calculated pK_a constants as compared to pure continuum models^{55,56} without making any further empirical adjustments besides the addition of explicit water molecules. Although Pliego and Riveros considered only a small number of compounds, their work suggests that for some ions, including explicit water molecules in the calculation is important for calculating pK_a . Adam carried out similar calculations¹³ for carboxylic acids and

phenols and found that the slope of the plot of experimental pK_a versus calculated acid dissociation free energy could be brought into much better agreement with experiment with the addition of explicit water molecules (again, explicit water molecules were added only to the ionic species). This work also suggests that accounting for explicit solvent effects may be necessary to make accurate predictions of pK_a .

An important challenge associated with adding explicit solvent molecules to implicit solvent calculations is that it is difficult to determine how many solvent molecules are required to reach convergence. The work of Pliego and Riveros⁵² and Adam¹³ illustrates this problem. In both studies different numbers of explicit solvent molecules (up to three in some cases) were used for different solutes (e.g., in the work of Adam, phenols required a single explicit water molecule to bring the calculated and experimental pK_a values into close agreement with one another, but carboxylic acids required two explicit water molecules). Thus, it is unclear how many explicit solvent molecules are required (if any) for compounds not included in their work.

The present article will examine what effect adding a single explicit solvent molecule to continuum solvation calculations has on the accuracy of pK_a predictions. We will also examine to what extent the addition of several explicit water molecules affects the accuracy of the prediction in cases where more than a single site exists on the conjugate base for strong hydrogen bonding interactions. For this, we will calculate the pK_a of bicarbonate using carbonate bound by up to three explicit water molecules.

2. Computing Acid Dissociation Free Energies from Solvation Free Energies

The standard-state free energy change associated with the following reaction, ΔG_{aq}^* ,



is related to the aqueous pK_a according to⁵⁷

$$pK_a = \frac{\Delta G_{\text{aq}}^*}{RT \ln(10)} \quad (2)$$

where R is the universal gas constant and T is the temperature. Using the thermodynamic cycle shown in Scheme 1, ΔG_{aq}^* may be expressed in terms of the aqueous solvation free energies of the acid AH and its conjugate base A^-

$$\Delta G_{\text{aq}}^*(\text{AH}) = \Delta G_{\text{gas}}^\circ(\text{AH}) + \Delta G_{\text{S}}^*(A^-) - \Delta G_{\text{S}}^*(\text{AH}) + \Delta G_{\text{S}}^*(\text{H}^+) + \Delta G^{\circ \rightarrow *}$$
 (3)

where $\Delta G_{\text{S}}^*(\text{AH})$ and $\Delta G_{\text{S}}^*(A^-)$ are the standard-state aqueous solvation free energies of AH and A^- , respectively, $\Delta G_{\text{S}}^*(\text{H}^+)$ is the aqueous solvation free energy of H^+ , $\Delta G_{\text{gas}}^\circ(\text{AH})$ is the gas-phase acidity of AH, defined by

$$\Delta G_{\text{gas}}^\circ(\text{AH}) = G_{\text{gas}}^\circ(A^-) + G_{\text{gas}}^\circ(\text{H}^+) - G_{\text{gas}}^\circ(\text{AH})$$
 (4)

and $\Delta G^{\circ \rightarrow *}$ is the free energy change associated with moving from a standard state that uses a concentration of 1 atm in the gas phase and 1 mol/L in the aqueous phase (denoted by the superscript “ \circ ”), to a standard state that uses a concentration of 1 mol/L in both the gas and aqueous phases (denoted by the superscript “ $*$ ”). The ideal-gas law yields

$$\Delta G^{\circ \rightarrow *} = RT \ln(24.46)$$
 (5)

At 298 K, $\Delta G^{\circ \rightarrow *}$ is 1.89 kcal/mol.

The acid dissociation free energy of AH in the aqueous phase may also be expressed in terms of the aqueous solvation free energy of the acid AH and its monohydrated conjugate base, $\text{H}_2\text{O} \cdot A^-$, using the thermodynamic cycle shown in Scheme 2

$$\Delta G_{\text{aq}}^*(\text{AH}) = \Delta G_{\text{gas}}^\circ(\text{H}_2\text{O} \cdot A^-) + \Delta G_{\text{S}}^*(\text{H}_2\text{O} \cdot A^-) - \Delta G_{\text{S}}^*(\text{AH}) - \Delta G_{\text{S}}^*(\text{H}_2\text{O}) + \Delta G_{\text{S}}^*(\text{H}^+)$$
 (6)

The gas-phase free energy $\Delta G_{\text{gas}}^\circ(\text{H}_2\text{O} \cdot A^-)$ in the above equation is equal to the sum of the gas-phase acidity of AH, $\Delta G_{\text{gas}}^\circ(\text{AH})$, and the gas-phase binding free energy of $\text{H}_2\text{O} \cdot A^-$, where this latter free energy is equal to $G_{\text{gas}}^\circ(\text{H}_2\text{O} \cdot A^-) - G_{\text{gas}}^\circ(\text{H}_2\text{O}) - G_{\text{gas}}^\circ(A^-)$. Note that the standard-state correction to the free energy, $\Delta G^{\circ \rightarrow *}$, does not appear in eq 6 because in Scheme 2, the number of moles on the products and reactants side is equal.

One can also replace H^+ by H_3O^+ in the above equations, and in fact we have recommended doing this for many purposes. However, for calculating pK_a values we recommend using the experimental value for $\Delta G_S^*(H^+)$, as discussed in Section 3. We also note that that $\Delta G_S^*(H^+)$ drops out for the important problem of calculating substituent effects on pK_a values (i.e., differences in pK_a values).

3. Experimental Data

All experimental and calculated data are for a temperature of 298 K. When available, we used experimental data to calculate ΔG_{aq}^* in eqs 3 and 6. For all of the acids studied here, experimental values for the gas-phase acidity, $\Delta G_{gas}^\circ(AH)$, and the aqueous solvation free energy (for neutral species) are available,^{47,58-60} and these were used in eqs 3 and 6. For the majority of the solute-water clusters studied here (monohydrated anions), experimental values are available⁶¹ for the gas-phase binding free energy, and these were used in eq 6. For the aqueous solvation free energy of the proton, $\Delta G_S^*(H^+)$, we used Tissandier et al.'s value of -265.9 kcal/mol.⁶² Experimental pK_a data were taken from the compilation of Stewart;⁶³ pK_a data not available in this compilation were taken from these additional sources.⁶⁴⁻⁶⁶

4. Computational Methods

In cases where experimental values for the gas-phase binding free energies are not available, we calculated them at the at the B97-1⁶⁷/MG3S⁶⁸ level of theory, using the *Gaussian 03* program.⁶⁹ (For atoms of atomic number 14 or lower, the MG3S basis set is identical to the older 6-311+G(2df,2p) basis^{70,71}). Aqueous solvation free energies of unclustered anions were calculated at the SM6/MPW25/6-31+G(d,p) level, using MPW25/MIDI! optimized gas-phase geometries (in Pople-style notation, this calculation is denoted SM6/MPW25/6-31+G(d,p)//MPW25/MIDI!). Note that MPW25 is a shorter name for the *mPW1PW91* functional of Adamo and Barone, which is also called *mPW0*.⁷² The aqueous solvation free energies of

clustered anions were calculated at the SM6/MPW25/6-31+G(d,p)//B97-1/MG3S level of theory. The SM6 method is implemented in the SMXGAUSS,⁷³ HONDOPLUS,⁷⁴ and GAMESSPLUS⁷⁵ programs.

5. Results and Discussion

Shown in Table 1 are the acid dissociation free energies calculated using eq 3, as well as the experimental pK_a values, for all 57 acids studied in this work (the additional column in this table is discussed below). For those acid dissociation free energies calculated using eq 3, we used experimental values for $\Delta G_{\text{gas}}^{\circ}(\text{AH})$, $\Delta G_{\text{S}}^*(\text{AH})$, and $\Delta G_{\text{S}}^*(\text{H}^+)$, and calculated values for $\Delta G_{\text{S}}^*(\text{A}^-)$. A plot of the experimental pK_a against the acid dissociation free energies obtained using eq 3 for all 57 acids yields a regression equation with a slope of $0.71/RT\ln(10)$ ($r^2 = 0.76$). This plot is shown in Figure 1a. To see what effect the addition of a single explicit water molecule would have on calculated acid dissociation free energies, we added a single water molecule around some of the conjugate bases (anions), and then used the calculated aqueous solvation free energies of the resulting solute-water clusters, $\Delta G_{\text{S}}^*(\text{H}_2\text{O} \cdot \text{A}^-)$, along with experimental values for $\Delta G_{\text{gas}}^{\circ}(\text{H}_2\text{O} \cdot \text{A}^-)$ (when available), $\Delta G_{\text{S}}^*(\text{AH})$, $\Delta G_{\text{S}}^*(\text{H}_2\text{O})$, and $\Delta G_{\text{S}}^*(\text{H}^+)$, to calculate ΔG_{aq}^* according to eq 6. The criteria we used to decide to which of the anions to add an explicit water molecule are the same criteria that we used during the development of our SM6 model.⁴⁷ In particular, we added a single explicit water molecule to any anion containing three or fewer atoms and to any anion with one or more oxygen atoms bearing a more negative partial atomic charge than bare water solute. Of the 57 acids in Table 1, 20 of them satisfy one or both of these criteria. The acid dissociation free energies calculated using eq 6 for these 20 acids are listed in Table 1. Plotting the experimental pK_a values of all 57 acids against the acid dissociation free energies obtained using eq 6 (20 of the acids in Table 1) combined with the remaining acid dissociation free energies using eq 3 (37 of the 57 acids in Table 1), yields a regression equation with a slope of $0.87/RT\ln(10)$ ($r^2 = 0.86$), which is a substantial improvement compared to when eq 3 was used to calculate acid dissociation free energies for all 57 acids. This plot is shown in Figure 1b.

The results obtained here can be compared to several other recent results. Klamt et al.^{23a} used an implicit solvent model^{76,77} to calculate aqueous acid dissociation free energies for 64 different acids, without the addition of explicit water molecules. These workers found a highly significant correlation ($r^2 = 0.98$) between the experimental pK_a and calculated acid dissociation free energy. However, the resulting slope was only $0.58/RT \ln(10)$, which is significantly lower than the expected slope of $1/RT \ln(10)$. On the basis of their results, as well as the results from two other studies,^{14,17} Klamt et al.²³ drew the startling conclusion that the experimental pK_a scale must depend differently on the free energy of dissociation than is generally assumed. Klicic et al.¹⁷ used eq 3, along with an implicit solvent model, to calculate dissociation free energies for a diverse data set containing 121 acids. With the exception of the 18 sulfonamides included in Klicic et al.'s data set (which required a regression equation different from that of the remaining acids), there is a strong correlation between experimental pK_a and the calculated dissociation free energy for the remaining 103 acids ($r^2 = 0.97$). However, the slope is only $0.53/RT \ln(10)$, which is very close to the value obtained by Klamt et al. Chipman¹⁴ calculated acid dissociation free energies in water for 6 different acids (water, methanol, ethanol, phenol, formic acid, acetic acid, and benzoic acid) using an implicit solvent model. Plotting experimental pK_a versus the calculated dissociation free energies reported by Chipman yields a slope of $0.51/RT \ln(10)$ ($r^2 = 0.99$). Interestingly, Chipman also calculated dissociation free energies for the six acids above in the aprotic solvent DMSO using the same implicit model that he used for water, and a plot of the experimental pK_a versus the calculated dissociation free energies for five of the six acids (an experimental pK_a value in DMSO is not available for formic acid), yields a regression equation with a slope equal to $0.84/RT \ln(10)$ ($r^2 = 0.99$). On the basis of these results, Chipman concluded, very reasonably, that for water, implicit solvent models alone are unable to account for strong anion-water interactions in the first solvation shell and suggested that in such cases where these interactions are expected to be important, better results might be obtained by augmenting continuum solvent calculations with other complementary methods that are especially designed to account for specific short-range interactions.

Adam calculated aqueous dissociation free energies using two approaches.¹³ In the first approach, an implicit solvent model was applied to 40 carboxylic acids and 20 phenols. A plot of the experimental pK_a values against the calculated dissociation free energies for each of these two classes of compounds results in two regression equations with slopes of $0.44/RT\ln(10)$ ($r^2 = 0.86$) and $0.50/RT\ln(10)$ ($r^2 = 0.97$), for the carboxylic acids and phenols, respectively. In the second approach, Adam repeated the implicit solvent calculations, each time successively adding a single explicit water molecule to the conjugate bases (the neutrals were modeled without explicit water molecules added throughout) until good agreement was obtained between the calculated and experimental slope of the resulting regression equation. For the phenols, adding a single water molecule to each of the anions results in a regression equation with a slope of $0.88/RT\ln(10)$ ($r^2 = 0.93$). For the carboxylic acids, a slope of $0.93/RT\ln(10)$ ($r^2 = 0.81$) was obtained, but only after two explicit water molecules were added.

On the basis of the results presented here, as well as the results obtained by several other workers, we can make several conclusions regarding the calculation of acid dissociation constants. First, although several other studies^{14,17,23} have shown that some implicit solvent models, without being augmented with explicit water molecules, can lead to very good predictions of the ordering of pK_a constants because there is a strong correlation between the calculated dissociation free energy and the experimental pK_a , the unusually low slopes of the resulting regression equations suggest that significant systematic errors are present in the underlying methods used to develop them.

Second, for some anions, an implicit model alone is unable to account for strong solute-solvent interactions in the first solvation shell, and for these types of solutes, treating one water molecule explicitly significantly increases the absolute accuracy of the calculated acid dissociation free energy. It is encouraging that a simple approach in which only a single water molecule is added to the calculation increases the accuracy because it avoids the problems⁴² introduced by the introduction of larger numbers of explicit solvent molecules (e.g., conformational sampling problems, computational cost). Furthermore, the above results are encouraging because, as pointed out in an earlier paper,⁴⁷ adding a small number of explicit water molecules to continuum

solvation calculations is much more straightforward than making large adjustments to the boundary between the solute and the continuum solvent (i.e. making large adjustments in the values used for the empirical atomic radii), using atom-typed radii,^{34,78-82} or using charge-dependent molecular cavities.⁸³⁻⁸⁹ Finally, applying this strategy yields a regression equation between the experimental pK_a and calculated acid dissociation free energy that obeys, reasonably well, the firmly founded thermodynamic relationship⁵⁷ between the dissociation free energy and pK_a .

Ideally we would give a definite prescription for cases in which a specific water molecule should be added to implicit solvent model calculations. However, it is not possible to do this in a way that covers the great diversity of possible cases that occur in applications. One prescription would be to add an explicit water whenever one wants to improve the accuracy, because adding an explicit water should almost always improve the accuracy when the effect is large, but it is relatively safe because it cannot make the accuracy much worse when the effect is small. The reason that adding explicit water might not always make the calculation more accurate is that the free energies of cavitation and dispersion and changes in the solute's translational, vibrational, and rotational free energy all change, and the parametrized surface tensions and the explicit solute-water and (possibly) water-water interactions are not necessarily accurate enough to account for small changes quantitatively. As an example of how large errors are reduced, we note that the pK_a value calculated here for phenol, whose conjugate base does not fall under one of the criteria explained above for adding explicit water, is 12.0. Adding a single explicit water molecule to phenoxide and using eq 6 leads to a calculated pK_a of 10.8, which is in better agreement with the experimental value of 10.0. We should also point out that although we obtain better results in most cases where we do add explicit water, for many of the acids studied here, we obtain reasonably good results without the explicit water. As an example of a case where excellent results are obtained without explicit water, so that the results cannot improve when water is added, we consider the calculated pK_a for ethanethiol, whose conjugate base also does not fall under either of the criteria that we used above for adding explicit water; this pK_a is in perfect agreement with the experimental pK_a without addition of

explicit water. Naturally, the agreement between the calculated and experimental pK_a does diminish when an explicit water molecule is added to the calculation, but only by a moderate amount (1.1 pK_a units). Despite cases such as this, the results above show that when one considers a statistically meaningful set of cases, addition of a single explicit water molecule improves the accuracy of calculated pK_a .

The next obvious question is: what effect does adding more than a single explicit water have on the accuracy of SM6? This question becomes especially important when one considers divalent ions, for which several sites may be available for strong solute-solvent hydrogen bonding interactions. To examine this more closely, we calculated the pK_a of the bicarbonate anion (HCO_3^-) using CO_3^{2-} bound by up to three water molecules. For comparative purposes, we also calculated the pK_a of HCO_3^- using the thermodynamic cycle shown in Scheme 1 (i.e., no bound waters):

$$pK_a(\text{HCO}_3^-) = (\Delta G_{\text{gas}}^\circ(\text{HCO}_3^-) + \Delta G_{\text{S}}^*(\text{CO}_3^{2-}) - \Delta G_{\text{S}}^*(\text{HCO}_3^-) + \Delta G_{\text{S}}^*(\text{H}^+) + \Delta G^{\circ \rightarrow *}) / 2.303 \log RT \quad (7)$$

where $\Delta G_{\text{gas}}^\circ(\text{HCO}_3^-)$ is the gas-phase acidity of HCO_3^- , and $\Delta G_{\text{S}}^*(\text{CO}_3^{2-})$ and $\Delta G_{\text{S}}^*(\text{HCO}_3^-)$ are the aqueous solvation free energies of CO_3^{2-} and HCO_3^- , respectively. The value that we used for $\Delta G_{\text{S}}^*(\text{HCO}_3^-)$ in eq 7 was obtained using the thermodynamic cycle shown in Scheme 1. For this, we used a theoretical⁹⁰ value of 332.2 kcal/mol for the gas-phase acidity of carbonic acid, (H_2CO_3), an experimental⁹¹ value of 6.4 for the pK_a of H_2CO_3 , the experimental⁶² value for the solvation free energy of the proton, and a calculated (SM6/MPW25/6-31+G(d,p)//MPW25/MIDI!) value for the aqueous solvation free energy of H_2CO_3 to arrive at a $\Delta G_{\text{S}}^*(\text{HCO}_3^-)$ value of -69 kcal/mol. The remaining free energy that is needed to calculate the pK_a of HCO_3^- with eq 7 is the aqueous solvation free energy of CO_3^{2-} , for which the SM6/MPW25/6-31+G(d,p)//B97-1/MG3S level of theory gives a value of -292 kcal/mol. Substituting the above values into eq 7 leads to a calculated pK_a of -6, which is in poor agreement with the experimental⁹² value of 10.3. This poor agreement is expected for unclustered calculations involving systems with such a large charge to volume ratio.

For pK_a calculations based on carbonate-water clusters we applied the thermodynamic cycle shown in

Scheme 2

$$pK_a(\text{HCO}_3^-) = (\Delta G_{\text{gas}}^\circ(\text{HCO}_3^-) + \Delta G_{\text{B.E.}}^\circ[(\text{H}_2\text{O})_n\text{CO}_3^{2-}] + \Delta G_{\text{S}}^*[(\text{H}_2\text{O})_n\text{CO}_3^{2-}] - \Delta G_{\text{S}}^*(\text{HCO}_3^-) - n\Delta G_{\text{S}}^*(\text{H}_2\text{O}) + \Delta G_{\text{S}}^*(\text{H}^+) - (n-1)\Delta G^{\circ \rightarrow *}) / 2.303 \log RT \quad (8)$$

where $\Delta G_{\text{gas}}^\circ(\text{HCO}_3^-)$ is the gas-phase acidity of HCO_3^- , $\Delta G_{\text{B.E.}}^\circ[(\text{H}_2\text{O})_n\text{CO}_3^{2-}]$ is the free energy associated with binding n water molecules to CO_3^{2-} in the gas phase, which is equal to

$G_{\text{gas}}^\circ[(\text{H}_2\text{O})_n\text{CO}_3^{2-}] - G_{\text{gas}}^\circ(\text{CO}_3^{2-}) - nG_{\text{gas}}^\circ(\text{H}_2\text{O})$, and $\Delta G_{\text{S}}^*[(\text{H}_2\text{O})_n\text{CO}_3^{2-}]$ is the aqueous solvation free energy of the resulting carbonate-water cluster. For $\Delta G_{\text{gas}}^\circ(\text{HCO}_3^-)$, we used a calculated value of 479 kcal/mol obtained at the G3⁹³ level of theory. This value is in perfect agreement with the value recently reported by Gao and co-workers⁴¹ for the same level of theory. Gas-phase binding energies of the three clusters were calculated at the B97-1/MG3S level of theory. Combining these values with the $\Delta G_{\text{S}}^*(\text{HCO}_3^-)$ value from above, experimental values for $\Delta G_{\text{S}}^*(\text{H}_2\text{O})$ and $\Delta G_{\text{S}}^*(\text{H}^+)$, and calculated (SM6/MPW25/6-31+G(d,p)//B97-1/MG3S) values for $\Delta G_{\text{S}}^*[(\text{H}_2\text{O})_n\text{CO}_3^{2-}]$, leads to calculated pK_a values of 0, 3, and 7 for $n = 1, 2$, and 3, respectively. These results are summarized in Table 2, along with the results obtained using the bare carbonate dianion. The structures of the three $(\text{H}_2\text{O})_n\text{CO}_3^{2-}$ clusters that were considered here are shown in Figure 2.

For SM6, the data in Table 2 are consistent with the statement above that adding explicit waters should almost always improve the results when the effect is large. Table 2 also shows that the accuracy of the pK_a prediction increases in a stepwise fashion as explicit waters are added. Finally, for some of the acids in Table 1, the above results suggest that the accuracy of the pK_a prediction can further be improved by the addition of more than one explicit water molecule to the conjugate base.

Finally, it is worth comparing the above results to those obtained using two other continuum solvation models: SM5.43R^{94,95} and the dielectric version^{55,96,97} of PCM (DPCM) with the UAHF method for assigning

atomic radii⁷⁹ as implemented in *Gaussian 98*⁹⁸ (we will refer to this second model as DPCM/98). For this, we calculated the pK_a of HCO_3^- using $\Delta G_S^*[(\text{H}_2\text{O})_n\text{CO}_3^{2-}]$ values (where $n = 0, 1, 2,$ or 3) calculated with SM5.43R and DPCM/98. The results of these calculations are summarized in Table 2. We chose the SM5.43R model for comparison against because it is the most recent previous version of our series of continuum solvation models. The older *Gaussian 98* implementation of DPCM was chosen because it was recently shown⁴⁷ to perform better than several other implementations of PCM for calculating aqueous solvation free energies of ionic solutes. The data in Table 2 show for DPCM/98, the closest agreement between the calculated and experimental pK_a of HCO_3^- is for $n = 0$, and the accuracy of the calculated pK_a *decreases* significantly as explicit water molecules are added to the calculation. For SM5.43R the best agreement is for $n = 1$, although adding more than one water molecule results in decreased accuracy.

6. Conclusions

The present article and our previous work⁴⁷ show that in many cases when an accurate implicit solvent model is used, the addition of explicit solvent molecules is not necessarily required for obtaining reasonable estimates of the pK_a . However, when strong specific solute-solvent hydrogen bonding interactions are expected to play an important role in the aqueous phase, adding a *single* explicit water molecule to the calculation significantly improves the accuracy of the pK_a prediction. This brings the slope of the pK_a scale into reasonable agreement with the experimental slope, and, despite the recent claims made by Klamt et al.,²³ should eliminate any doubts as to the correctness of the standard thermodynamic treatment⁵⁷ of pK_a .

To see what effect adding multiple explicit waters has on the accuracy of calculated pK_a values in cases where several sites are available for strong solute-solvent hydrogen-bonding interactions, we calculated the pK_a of HCO_3^- using carbonate-water clusters bound by up to three explicit water molecules. It was shown that the accuracy of the prediction significantly improved with each addition of an explicit molecule—the calculated and experimental pK_a of HCO_3^- are in better agreement when three explicit water molecules are added to the

conjugate base (CO_3^{2-}). When two other continuum models, SM5.43R and PCM, are used instead of SM6, the accuracy of the calculated pK_a for HCO_3^- significantly worsens when three explicit water molecules are added.

Acknowledgment. This work was supported by the NIH training grant for Neuro-physical-computational Sciences, by the U.S. Army Research Office under Multidisciplinary Research Program of the University Research Initiative (MURI) through grant number DAAD19-02-1-0176, by the Minnesota Partnership for Biotechnology and Medical Genomics, by the National Science Foundation (CHE02-03446 and CHEO3-49122), and by the Office of Naval Research under grant no. N 00014-05-01-0538.

References

- (1) Hammett, L. P. *J. Am. Chem. Soc.* **1937**, *59*, 96.
- (2) Schuurman, G.; Cossi, M.; Barone, V.; Tomasi, J. *J. Phys. Chem. A* **1998**, *102*, 6706.
- (3) da Silva, C. O.; da Silva, E. C.; Nascimento, M. A. C. *J. Phys. Chem. A* **1999**, *103*, 11194.
- (4) Vogel, H. J.; Juffer, A. H. *Theor. Chem. Acct.* **1999**, *101*, 159.
- (5) Chen, I.-J.; MacKerell Jr, A. D. *Theor. Chem. Acct.* **2000**, *103*, 483.
- (6) Silva, C. O.; da Silva, E. C.; Nascimento, M. A. C. *J. Phys. Chem. A* **2000**, *104*, 2402.
- (7) Topol, I. A.; Tawa, G. J.; Caldwell, R. A.; Eissenstat, M. A.; Burt, S. K. *J. Phys. Chem. A* **2000**, *104*, 9619.
- (8) Jang, Y. H.; Sowers, L. C.; Çagin, T.; Goddard III, W. A. *J. Phys. Chem. A* **2001**, *105*, 274.
- (9) Liptak, M. D.; Shields, G. C. *Int. J. Quantum Chem.* **2001**, *85*, 727.
- (10) Liptak, M. D.; Shields, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 7314.
- (11) Ohno, K.; Kamiya, N.; Asakawa, N.; Inoue, Y.; Sakurai, M. *Chem. Phys. Lett.* **2001**, *341*, 387.
- (12) Toth, A. M.; Liptak, M. D.; Phillips, D. L.; Shields, G. C. *J. Chem. Phys.* **2001**, *114*, 4595.
- (13) Adam, K. *J. Phys. Chem. A* **2002**, *106*, 11963.
- (14) Chipman, D. M. *J. Phys. Chem. A* **2002**, *106*, 7413.
- (15) Davies, J. E.; Doltsinis, N. L.; Kirby, A. J.; Roussev, C. D.; Sprik, M. *J. Am. Chem. Soc.* **2002**, *124*, 6594.

- (16) Feierberg, I.; Åqvist, J. *Theor. Chem. Acct.* **2002**, *108*, 71.
- (17) Klicić, J. J.; Friesner, R. A.; Liu, S.-Y.; Guida, W. C. *J. Phys. Chem. A* **2002**, *106*, 1327.
- (18) Li, H.; Hains, A. W.; Everts, J. E.; Robertson, A. D.; Jensen, J. H. *J. Phys. Chem. B* **2002**, *106*, 3486.
- (19) Liptak, M. D.; Gross, K. C.; Seybold, P. G.; Feldgus, S.; Shields, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 6421.
- (20) Lopez, X.; Schaefer, M.; Dejaegere, A.; Karplus, M. *J. Am. Chem. Soc.* **2002**, *124*, 5010.
- (21) Satchell, J. F.; Smith, B. J. *Phys. Chem. Chem. Phys.* **2002**, *4*, 4314.
- (22) Jang, Y. H.; Goddard III, W. A.; Noyes, K. T.; Sowers, L. C.; Hwang, S.; Chung, D. S. *J. Phys. Chem. B* **2003**, *107*, 344.
- (23) (a) Klamt, A.; Eckert, F.; Diedenhofen, M.; Beck, M. E. *J. Phys. Chem. A* **2003**, *107*, 9380 (b) Eckert, F.; Klamt, A. *J. Comput. Chem.* **2006**, *27*, 11.
- (24) Li, G.; Cui, Q. *J. Phys. Chem. B* **2003**, *107*, 14521.
- (25) Mujika, J. I.; Mercero, J. M.; Lopez, X. *J. Phys. Chem. A* **2003**, *107*, 6099.
- (26) Almerindo, G. I.; Tondo, D. W.; Pliego_Jr, J. R. *J. Phys. Chem. A* **2004**, *108*, 166.
- (27) Barone, V.; Improta, R.; Rega, N. *Theor. Chem. Acct.* **2004**, *111*, 237.
- (28) da Silva, C. O.; Barbosa, A. G. H.; da Silva, E. T.; da Silva, E. L. L.; Nascimento, M. A. C. *Theor. Chem. Acct.* **2004**, *111*, 231.
- (29) Fu, Y.; Liu, L.; Li, R.-Q.; Liu, R.; Guo, Q.-X. *J. Am. Chem. Soc.* **2004**, *126*, 814.
- (30) Namazian, M.; Halvani, S.; Noorbala, M. R. *J. Mol. Struct. (THEOCHEM)* **2004**, *711*, 13.
- (31) Quenneville, J.; Popovic, D. M.; Stuchebrukhov, A. A. *J. Phys. Chem. B* **2004**, *108*, 18383.
- (32) Schmidt am Busch, M.; Knapp, E. W. *ChemPhysChem* **2004**, *5*, 1513.
- (33) Simonson, T.; Carlsson, J.; Case, D. A. *J. Am. Chem. Soc.* **2004**, *126*, 4167.
- (34) Curutchet, C.; Bidon-Chanal, A.; Soteras, I.; Orozco, M.; Luque, F. J. *J. Phys. Chem. B* **2005**, *109*, 3565.
- (35) Han, J.; Deming, R. L.; Tao, F.-M. *J. Phys. Chem. A* **2005**, *109*, 1159.
- (36) Jensen, J. H.; Li, H.; Robertson, A. D.; Molina, P. A. *J. Phys. Chem. A* **2005**, *109*, 6634.

- (37) Kaminski, G. A. *J. Phys. Chem. B* **2005**, *109*, 5884.
- (38) Popovic, D. M.; Quenneville, J.; Stucherbrukhov, A. A. *J. Phys. Chem. B* **2005**, *109*, 3616.
- (39) Takano, Y.; Houk, K. N. *J. Chem. Theory Comput.* **2005**, *1*, 70.
- (40) Wang, X.; Li, S.; Jiang, Y. *J. Phys. Chem. A* **2005**, *109*, 10776.
- (41) Gao, D.; Svoronos, P.; Wong, P. K.; Maddalena, D.; Hwang, J.; Walker, H. *J. Phys. Chem. A* **2005**, *109*, 10776.
- (42) Cramer, C. J.; Truhlar, D. G. *Chem. Rev.* **1999**, *99*, 2161.
- (43) Tomasi, J.; Mennucci, B.; Cammi, R. *Chem. Rev.* **2005**, *105*, 2999.
- (44) Laria, D.; Kapral, R.; Estrin, D.; Cicott, G. J. *J. Chem. Phys.* **1996**, *104*, 6560.
- (45) Cramer, C. J.; Truhlar, D. G. *J. Am. Chem. Soc.* **1991**, *113*, 8305.
- (46) Chambers, C. C.; Hawkins, G. D.; Cramer, C. J.; Truhlar, D. G. *J. Phys. Chem.* **1996**, *100*, 16385.
- (47) Kelly, C. P.; Cramer, C. J.; Truhlar, D. G. *J. Chem. Theory Comput.* **2005**, *1*, 1133.
- (48) Claverie, P.; Daudey, J. P.; Langlet, J.; Pullman, B.; Piazzola, D.; Huron, M. J. *J. Phys. Chem.* **1978**, *82*, 405.
- (49) Sánchez-Marcos, E.; Pappalardo, R. R.; Rinaldi, D. *J. Phys. Chem.* **1991**, *95*, 8928.
- (50) Tuñoñ, I.; Rinaldi, J.-L.; Ruiz-López, M. F.; Rivail, J. L. *J. Phys. Chem.* **1995**, *99*, 3798.
- (51) Pliego Jr., J. R.; Riveros, J. M. *J. Phys. Chem. A* **2001**, *105*, 7241.
- (52) Pliego Jr., J. R.; Riveros, J. M. *J. Phys. Chem. A* **2002**, *106*, 7434.
- (53) Karelson, M.; Zerner, M. C. *J. Am. Chem. Soc.* **1990**, *112*, 9405.
- (54) Fox, T.; Rösch, N. *Chem. Phys. Lett.* **1992**, *191*, 33.
- (55) Cossi, M.; Barone, V.; Cammi, R.; Tomasi, J. *Chem. Phys. Lett.* **1996**, *255*, 327.
- (56) Li, J.; Zhu, T.; Hawkins, G. D.; Winget, P.; Liotard, D. A.; Cramer, C. J.; Truhlar, D. G. *Theor. Chem. Acc.* **1999**, *103*, 9.
- (57) Lewis, G. N.; Randall, M.; Pitzer, K. S.; Brewer, L. *Thermodynamics*, 2nd ed.; McGraw-Hill: New York, 1961.

- (58) *Physical/Chemical Property Database (PHYSPROP)*; SRC Environmental Science Center: Syracuse, NY, 1994.
- (59) O'Sullivan, D. W.; Lee, M.; Noone, B. C.; Heikes, B. G. *J. Phys. Chem.* **1996**, *100*, 3241.
- (60) Régimbal, J. M.; Mozurkewich, M. *J. Phys. Chem. A* **1997**, *101*, 8822.
- (61) Meot-Ner, M. M.; Lias, S. G. Binding Energies Between Ions and Molecules, and the Thermochemistry of Cluster Ions. In *NIST Chemistry WebBook, NIST Standard Reference Database Number 69*; Linstrom, P. J., Mallard, W. G., Eds.; National Institute of Standards and Technology: Gaithersburg, MD, March 2003.
- (62) (a) Tissandier, M. D.; Cowen, K. A.; Feng, W. Y.; Gundlach, E.; Cohen, M. J.; Earhart, A. D.; Coe, J. V. *J. Phys. Chem. A* **1998**, *102*, 7784 (b) Camaioni, D. M.; Schwerdtfeger, C. A. *J. Phys. Chem. A* **2005**, *109*, 10795.
- (63) Stewart, R. *The Proton: Applications to Organic Chemistry*; Academic: New York, 1985.
- (64) Perrin, D. D. *Ionisation Constants of Inorganic Acids and Bases in Aqueous Solution*; Pergamon: New York, 1982.
- (65) Albert, A.; Serjeant, E. P. *The Determination of Ionization Constants: A Laboratory Manual*; Chapman and Hall: New York, 1984.
- (66) Kresge, A. J.; Pruszynski, P.; Stang, P. T.; Williamson, B. J. *J. Org. Chem.* **1991**, *56*, 4808.
- (67) Hamprecht, F. A.; Cohen, A. J.; Tozer, D. J.; Handy, N. C. *J. Chem. Phys.* **1998**, *109*, 6264.
- (68) Lynch, B. J.; Zhao, Y.; Truhlar, D. G. *J. Phys. Chem. A* **2003**, *107*, 1384.
- (69) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery Jr., J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalamani, G.; Rega, N.; Petersson, G. A.; Nagatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A.

G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. Gaussian 03, Revision C.01 Pittsburgh PA, 2003.

(70) Krishnan, R.; Binkley, J. S.; Seeger, R.; Pople, J. A. *J. Chem. Phys.* **1980**, *72*, 650.

(71) Frisch, M. J.; Pople, J. A.; Binkley, J. S. *J. Chem. Phys.* **1984**, *80*, 3265.

(72) Adamo, C.; Barone, V. *J. Chem. Phys.* **1998**, *108*, 664.

(73) Thompson, D. J.; Kelly, C. P.; Lynch, B. J.; Xidos, J. D.; Li, J.; Hawkins, G. D.; Zhu, T.; Volobuev, Y.; Dupuis, M.; Rinaldi, D.; Liotard, D. A.; Cramer, C. J.; Truhlar, D. G. SMXGAUSS-version 3.2, University of Minnesota, Minneapolis, MN 55455, 2005.

(74) Nakamura, H.; Kelly, C. P.; Xidos, J. D.; Thompson, D. J.; Li, J.; Hawkins, G. D.; Zhu, T.; Lynch, B. J.; Volobuev, Y.; Rinaldi, D.; Liotard, D. A.; Cramer, C. J.; Truhlar, D. G. HONDOPLUS-version 4.9 University of Minnesota, Minneapolis, 2005, based on HONDO v. 99.6.

(75) Pu, J.; Kelly, C. P.; Thompson, D. J.; Xidos, J. D.; Li, J.; Zhu, T.; Hawkins, G. D.; Chuang, Y.-Y.; Fast, P. L.; Lynch, B. J.; Liotard, D. A.; Rinaldi, D.; Gao, J.; Cramer, C. J.; Truhlar, D. G. GAMESSPLUS-version 4.7, University of Minnesota, Minneapolis, 2005, based on the General Atomic and Molecular Electronic Structure System (GAMESS) as described in Schmidt, M. W.; Baldridge, K. K.; Boatz, J. A.; Elbert, S. T.; Gordon, M. S.; Jensen, J. H.; Koseki, S.; Matsunaga, N.; Nguyen, K. A.; Su, S. J.; Windus, T. L.; Dupuis, M.; Montgomery, J. A. *J. Comput. Chem.* **1993**, *14*, 1347-1363.

(76) Klamt, A. *J. Phys. Chem.* **1995**, *99*, 2224.

(77) Klamt, A.; Jonas, V.; Bürger, T.; Lohrenz, J. C. W. *J. Phys. Chem.* **1998**, *102*, 5074.

(78) Marten, B.; Kim, K.; Cortis, C.; Friesner, R.; Murphy, R. B.; Ringnalda, M. N.; Sitkoff, D.; Honig, B. *J. Phys. Chem.* **1996**, *100*, 11775.

(79) Barone, V.; Cossi, M.; Tomasi, J. *J. Chem. Phys.* **1997**, *107*, 3210.

(80) Nina, M.; Beglov, D.; Roux, B. *J. Phys. Chem. B* **1997**, *101*, 5239.

- (81) Banavali, N. K.; Roux, B. *J. Phys. Chem. B* **2002**, *106*, 11026.
- (82) Swanson, J. M. J.; Adcock, S. A.; McCammon, J. A. *J. Chem. Theory. Comput.* **2005**, *1*, 484.
- (83) Aguilar, M. A.; Martin, M. A.; Tolosa, S.; Olivares del Valle, F. J. *J. Mol. Struct. (THEOCHEM)* **1988**, *166*, 313.
- (84) Aguilar, M. A.; Olivares del Valle, F. J. *Chem. Phys.* **1989**, *129*, 439.
- (85) Gonçalves, P. F. B.; Livotto, P. R. *Chem. Phys. Lett.* **1999**, *304*, 438.
- (86) Olivares del Valle, F. J.; Aguilar, M. A.; Contador, J. C. *Chem. Phys.* **1993**, *170*, 161.
- (87) Rick, S. W.; Berne, B. J. *J. Am. Chem. Soc.* **1994**, *116*, 3949.
- (88) Takahashi, O.; Sawahata, H.; Ogawa, Y.; Kikuchi, O. *J. Mol. Struct. (THEOCHEM)* **1997**, *393*, 141.
- (89) Camaioni, D. M.; Dupuis, M.; Bentley, J. *J. Phys. Chem. A* **2003**, *107*, 5778.
- (90) Alexeev, Y.; Windus, T. L.; Zhan, C.-G.; Dixon, D. A. *Int. J. Quantum Chem.* **2005**, *102*, 775.
- (91) Schwarzenbach, G. *Helv. Chim. Acta* **1957**, *40*, 907.
- (92) Lide, D. R. *CRC Handbook of Chemistry and Physics, 80th edn.*; CRC Press: Boca Raton, 2000.
- (93) Curtiss, L. A.; Raghavachari, K.; Redfern, P.; Rassolov, V.; Pople, J. A. *J. Chem. Phys.* **1998**, *109*, 7764.
- (94) Thompson, J. D.; Cramer, C. J.; Truhlar, D. G. *Theor. Chem. Acc.* **2005**, *113*, 107.
- (95) Thompson, J. D.; Cramer, C. J.; Truhlar, D. G. *J. Phys. Chem. A* **2004**, *108*, 6532.
- (96) Miertus, S.; Scrocco, E.; Tomasi, J. *J. Chem. Phys.* **1981**, *55*, 117.
- (97) Miertus, S.; Tomasi, J. *Chem. Phys.* **1982**, *65*, 239.
- (98) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; Montgomery Jr., J. A.; Stratmann, R. E.; Burant, J. C.; Dapprich, S.; Millam, J. M.; Daniels, A. D.; Kudin, K. N.; Strain, M. C.; Farkas, O.; Tomasi, J.; Barone, V.; Cossi, M.; Cammi, R.; Mennucci, B.; Pomelli, C.; Adamo, C.; Clifford, S.; Ochterski, J. W.; Petersson, G. A.; Ayala, P. Y.; Cui, Q.; Morokuma, K.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Cioslowski, J.; Ortiz, J. V.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Gonzalez, C.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong,

M. W.; Andres, J. L.; Head-Gordon, M.; Replogle, E. S.; Pople, J. A. Gaussian 98, Revision A.11 Pittsburgh, PA, 1998.

TABLE 1: Calculated Dissociation Free Energies (SM6) Used to Develop the Correlation $pK_a = m \Delta G_{aq}^* / RT \ln(10) + b$.

AH	ΔG_{aq}^* (eq 3) ^a Implicit Solvent	ΔG_{aq}^* (eq 6) ^b Implicit + Explicit Solvent	Expt. pK_a
trifluoroacetic acid	-3.2		0.5
dichloroacetic acid	0.5		1.4
pyruvic acid	0.5		2.5
chloroacetic acid	5.7		2.9
formic acid	3.0		3.8
benzoic acid	3.8		4.2
acrylic acid	4.0		4.3
hydroperoxyl radical	-4.4	3.8	4.7
acetic acid	7.1		4.8
propanoic acid	8.9		4.9
hexanoic acid	9.4		4.9
thiophenol	6.6		6.6
hydrogen sulfide	1.8	3.4	7.0
4-nitrophenol	11.1		7.1
2-nitrophenol	4.5		7.2
3-nitrophenol	14.3		8.4
2-chlorophenol	11.8		8.5
hydrogen cyanide	-3.4	1.1	9.2
1,1,1,3,3,3-hexafluoropropan-2-ol	19.0	17.8	9.3
3-hydroxyphenol	16.3		9.3
4-chlorophenol	15.1		9.4
4-hydroxyphenol	19.5		9.9
phenol	16.3		10.0
3-methylphenol	15.8		10.1
nitromethane	15.8		10.2
4-methylphenol	17.2		10.3
cyanamide	10.1		10.3
methanethiol	12.1		10.3
2-methylphenol	16.9		10.3
ethanethiol	14.4		10.6
1-propanethiol	14.7		10.7
methyl hydroperoxide	20.7	21.1	11.5
hydrogen peroxide	14.8	23.1	11.7
ethyl hydroperoxide	18.7	18.1	11.8
2,2,2-trifluoroethanol	24.4	19.7	12.4
2-methoxyethanol	29.7	23.6	14.8
acetamide	22.2		15.1
benzyl alcohol	32.0	25.7	15.4
1,2-ethanediol	21.6	15.7	15.4
methanol	27.8	21.8	15.5

allyl alcohol	28.6	22.2	15.5
water	9.7	17.3	15.7
ethanol	30.2	24.0	15.9
1-propanol	30.0	22.7	16.1
acetaldehyde	20.8		16.5
isopropanol	31.8	26.0	17.1
2-butanol	35.2	28.7	17.6
4-nitroaniline	25.3		18.2
acetone	27.1		19.0
t-butanol	34.6	27.5	19.2
3-pentanone	32.1		19.9
acetylene	19.2	23.3	21.7
diphenylamine	27.1		22.4
chloroform	33.3		24.0
acetonitrile	32.1		25.0
aniline	33.2		27.7
dimethyl sulfoxide	40.6		33.0
<i>X/Y</i> ^c	57/0	37/20	
<i>m</i>	0.71	0.87	
<i>b</i>	3.10	1.39	
<i>r</i> ²	0.76	0.86	

^aAqueous dissociation free energy, calculated using experimental values for $\Delta G_{\text{gas}}^{\circ}(\text{A}^-)$, $\Delta G_{\text{S}}^*(\text{AH})$, and $\Delta G_{\text{S}}^*(\text{H}^+)$, and calculated values for $\Delta G_{\text{S}}^*(\text{A}^-)$, in eq 3.

^bAqueous dissociation free energy, calculated using experimental values for $\Delta G_{\text{gas}}^{\circ}(\text{H}_2\text{O} \cdot \text{A}^-)$, $\Delta G_{\text{S}}^*(\text{AH})$, $\Delta G_{\text{S}}^*(\text{H}_2\text{O})$, and $\Delta G_{\text{S}}^*(\text{H}^+)$, and calculated values for $\Delta G_{\text{S}}^*(\text{H}_2\text{O} \cdot \text{A}^-)$, in eq 6. ^c*X* is the number of ΔG_{aq}^* values calculated using eq 3 that were used to develop the correlation between experimental $\text{p}K_{\text{a}}$ and ΔG_{aq}^* ; *Y* is the number of ΔG_{aq}^* values calculated using eq 6 that were used to develop the correlation between experimental $\text{p}K_{\text{a}}$ and ΔG_{aq}^* .

TABLE 2: Gas-Phase Binding Free Energies of $(\text{H}_2\text{O})_n\text{CO}_3^{2-}$ (kcal/mol) and Calculated $\text{p}K_a$ Values^a of HCO_3^- Obtained without ($n = 0$) and with ($n > 0$) the Addition of Explicit Water Molecules to CO_3^{2-} , Using Different Continuum Solvation Models for Calculating the Aqueous Solvation Free Energy of $(\text{H}_2\text{O})_n\text{CO}_3^{2-}$. (Experimental $\text{p}K_a$ of $\text{HCO}_3^- = 10.3$).

n	$\Delta G_{\text{B.E.}}^\circ[(\text{H}_2\text{O})_n\text{CO}_3^{2-}]^b$	SM6 ^c	SM5.43R ^d	DPCM/98 ^e
0	0	-6 (16)	6 (4)	16 (6)
1	-28	0 (10)	11 (1)	18 (8)
2	-49	3 (7)	16 (6)	23 (13)
3	-67	7 (3)	20 (10)	35 (25)


^aAbsolute deviations relative to experiment are given in parentheses.

^bCalculated (B97-1/MG3S) values.


^cSM6/MPW25/6-31+G(d,p)//B97-1/MG3S

^dSM5.43R/MPW25/6-31+G(d,p)//B97-1/MG3S

^eDPCM/98/HF/6-31G(d)//B97-1/MG3S; UAHF atomic radii used.



Scheme 1



Scheme 2





Figure 1. Experimental pK_a plotted against the calculated acid dissociation free energy. The dashed line is the expected result obtained using eq 2 (slope = $1/RT\ln(10)$; intercept = 0). Solid circles indicate acid dissociation free energies calculated using eq 3 (no explicit water). (a) Open squares indicate acids whose dissociation free energies will be calculated using eq 6 (single explicit water molecule; see Figure 1b). (b) Open circles indicate acid dissociation energies calculated using eq 6 (single explicit water molecule added to the conjugate base).




Figure 2. Carbonate-water clusters considered in this study. Dashed lines represent hydrogen bonds.